arXiv:1401.0815v2 [math.CA] 12 Jan 2015

Additions to the formula lists in "Hypergeometric orthogonal polynomials and their q-analogues" by Koekoek, Lesky and Swarttouw

Tom H. Koornwinder

Abstract

This report gives a rather arbitrary choice of formulas for (q-)hypergeometric orthogonal polynomials which the author missed while consulting Chapters 9 and 14 in the book "Hypergeometric orthogonal polynomials and their q-analogues" by Koekoek, Lesky and Swarttouw. The systematics of these chapters will be followed here, in particular for the numbering of subsections and of references.

Introduction

This report contains some formulas about (q-)hypergeometric orthogonal polynomials which I missed but wanted to use while consulting Chapters 9 and 14 in the book

R. Koekoek, P. A. Lesky and R. F. Swarttouw, *Hypergeometric orthogonal polynomials and their q-analogues*, Springer-Verlag, 2010.

These chapters form together the (slightly extended) successor of the report

R. Koekoek and R. F. Swarttouw, *The Askey-scheme of hypergeometric orthogonal polynomials and its q-analogue*, Report 98-17, Faculty of Technical Mathematics and Informatics, Delft University of Technology, 1998; `http://aw.twi.tudelft.nl/~koekoek/askey/`.

Certainly these chapters give complete lists of formulas of special type, for instance orthogonality relations and three-term recurrence relations. But outside these narrow categories there are many other formulas for (q-)orthogonal polynomials which one wants to have available. Often one can find the desired formula in one of the standard references listed at the end of this report. Sometimes it is only available in a journal or a less common monograph. Just for my own comfort, I have brought together some of these formulas. This will possibly also be helpful for some other users.

Usually, any type of formula I give for a special class of polynomials, will suggest a similar formula for many other classes, but I have not aimed at completeness by filling in a formula of such type at all places. The resulting choice of formulas is rather arbitrary, just depending on the formulas which I happened to need or which raised my interest. For each formula I give a suitable reference or I sketch a proof. It is my intention to gradually extend this collection of formulas.

Conventions

The (x.y) and (x.y.z) type subsection numbers, the (x.y.z) type formula numbers, and the [x] type citation numbers refer to the book by Koekoek et al. The (x) type formula numbers refer to this manuscript and the [Kx] type citation numbers refer to citations which are not in the book. Some standard references like [DLMF] are given by special acronyms.

N is always a positive integer. Always assume n to be a nonnegative integer or, if N is present, to be in $\{0, 1, \ldots, N\}$. Throughout assume $0 < q < 1$.

For each family the coefficient of the term of highest degree of the orthogonal polynomial of degree n can be found in the book as the coefficient of $p_n(x)$ in the formula after the main formula under the heading "Normalized Recurrence Relation". If that main formula is numbered as (x.y.z) then I will refer to the second formula as (x.y.zb).

Contents

Introduction
Conventions
Generalities

9.1 Wilson
9.2 Racah
9.3 Continuous dual Hahn
9.4 Continuous Hahn
9.5 Hahn
9.6 Dual Hahn
9.7 Meixner-Pollaczek
9.8 Jacobi
    9.8.1 Gegenbauer / Ultraspherical
    9.8.2 Chebyshev
9.9 Pseudo Jacobi (or Routh-Romanovski)
9.10 Meixner
9.11 Krawtchouk
9.12 Laguerre
9.14 Charlier
9.15 Hermite

14.1 Askey-Wilson
14.2 q-Racah
14.3 Continuous dual q-Hahn
14.4 Continuous q-Hahn
14.5 Big q-Jacobi
14.7 Dual q-Hahn
14.8 Al-Salam-Chihara
14.10 Continuous q-Jacobi
    14.10.1 Continuous q-ultraspherical / Rogers
14.11 Big q-Laguerre
14.12 Little q-Jacobi
14.14 Quantum q-Krawtchouk
14.16 Affine q-Krawtchouk
14.17 Dual q-Krawtchouk
14.20 Little q-Laguerre
14.21 q-Laguerre
14.27 Stieltjes-Wigert
14.28 Discrete q-Hermite I
14.29 Discrete q-Hermite II

Standard references
References from Koekoek, Lesky & Swarttouw
Other references

Generalities

Criteria for uniqueness of orthogonality measure According to Shohat & Tamarkin [K24, p.50] orthonormal polynomials p_n have a unique orthogonality measure (up to positive constant factor) if for some $z \in \mathbb{C}$ we have

$$\sum_{n=0}^{\infty} |p_n(z)|^2 = \infty. \tag{1}$$

Also (see Shohat & Tamarkin [K24, p.59]), monic orthogonal polynomials p_n with three-term recurrence relation $xp_n(x) = p_{n+1}(x) + B_n p_n(x) + C_n p_{n-1}(x)$ (C_n necessarily positive) have a unique orthogonality measure if

$$\sum_{n=1}^{\infty} (C_n)^{-1/2} = \infty. \tag{2}$$

Furthermore, if orthogonal polynomials have an orthogonality measure with bounded support, then this is unique (see Chihara [146]).

Even orthogonality measure If $\{p_n\}$ is a system of orthogonal polynomials with respect to an even orthogonality measure which satisfies the three-term recurrence relation

$$xp_n(x) = A_n p_{n+1}(x) + C_n p_{n-1}(x)$$

then

$$\frac{p_{2n}(0)}{p_{2n-2}(0)} = -\frac{C_{2n-1}}{A_{2n-1}}\,. \tag{3}$$

Appell's bivariate hypergeometric function F_4 This is defined by

$$F_4(a,b;c,c';x,y) := \sum_{m,n=0}^{\infty} \frac{(a)_{m+n}(b)_{m+n}}{(c)_m (c')_n\, m!\, n!}\, x^m y^n \qquad (|x|^{\frac12} + |y|^{\frac12} < 1), \tag{4}$$

see [HTF1, 5.7(9), 5.7(44)] or [DLMF, (16.13.4)]]. There is the reduction formula

$$F_4\left(a,b;b,b;\frac{-x}{(1-x)(1-y)},\frac{-y}{(1-x)(1-y)}\right) = (1-x)^a(1-y)^a\ {}_2F_1\left(\begin{matrix}a, 1+a-b\\ b\end{matrix};xy\right),$$

see [HTF1, 5.10(7)]. When combined with the quadratic transformation [HTF1, 2.11(34)] (here $a-b-1$ should be replaced by $a-b+1$), see also [DLMF, (15.8.15)]], this yields

$$F_4\left(a,b;b,b;\frac{-x}{(1-x)(1-y)},\frac{-y}{(1-x)(1-y)}\right) = \left(\frac{(1-x)(1-y)}{1+xy}\right)^a\ {}_2F_1\left(\begin{matrix}\frac12 a, \frac12(a+1)\\ b\end{matrix};\frac{4xy}{(1+xy)^2}\right).$$

This can be rewritten as

$$F_4(a,b;b,b;x,y) = (1-x-y)^{-a}\ {}_2F_1\left(\begin{matrix}\frac12 a, \frac12(a+1)\\ b\end{matrix};\frac{4xy}{(1-x-y)^2}\right). \tag{5}$$

Note that, if $x, y \geq 0$ and $x^{\frac12} + y^{\frac12} < 1$, then $1-x-y > 0$ and $0 \leq \frac{4xy}{(1-x-y)^2} < 1$.

q-Hypergeometric series of base q^{-1} By [GR, Exercise 1.4(i)]:

$$ {}_r\phi_s\left(\begin{matrix}a_1,\dots,a_r\\ b_1,\dots b_s\end{matrix};q^{-1},z\right)= {}_{s+1}\phi_s\left(\begin{matrix}a_1^{-1},\dots a_r^{-1},0,\dots,0\\ b_1^{-1},\dots,b_s^{-1}\end{matrix};q,\frac{qa_1\dots a_rz}{b_1\dots b_s}\right) \tag{6}$$

for $r\le s+1$, $a_1,\dots,a_r,b_1,\dots,b_s\neq 0$. In the non-terminating case, for $0<q<1$, there is convergence if $|z|<b_1\dots b_s/(qa_1\dots a_r)$.

A transformation of a terminating ${}_2\phi_1$ By [GR, Exercise 1.15(i)] we have

$$ {}_2\phi_1\left(\begin{matrix}q^{-n},b\\ c\end{matrix};q,z\right)=(bz/(cq);q^{-1})_n\ {}_3\phi_2\left(\begin{matrix}q^{-n},c/b,0\\ c,cq/(bz)\end{matrix};q,q\right). \tag{7}$$

Very-well-poised q-hypergeometric series The notation of [GR, (2.1.11)] will be followed:

$$ {}_{r+1}W_r(a_1;a_4,a_5,\dots,a_{r+1};q,z):= {}_{r+1}\phi_r\left(\begin{matrix}a_1,qa_1^{\frac12},-qa_1^{\frac12},a_4,\dots,a_{r+1}\\ a_1^{\frac12},-a_1^{\frac12},qa_1/a_4,\dots,qa_1/a_{r+1}\end{matrix};q,z\right). \tag{8}$$

Theta function The notation of [GR, (11.2.1)] will be followed:

$$ \theta(x;q):=(x,q/x;q)_\infty,\qquad \theta(x_1,\dots,x_m;q):=\theta(x_1;q)\dots\theta(x_m;q). \tag{9}$$

9.1 Wilson

Symmetry The Wilson polynomial $W_n(y;a,b,c,d)$ is symmetric in a,b,c,d.
This follows from the orthogonality relation (9.1.2) together with the value of its coefficient of y^n given in (9.1.5b). Alternatively, combine (9.1.1) with [AAR, Theorem 3.1.1].
As a consequence, it is sufficient to give generating function (9.1.12). Then the generating functions (9.1.13), (9.1.14) will follow by symmetry in the parameters.

Hypergeometric representation In addition to (9.1.1) we have (see [513, (2.2)]):

$$\begin{aligned} W_n(x^2;a,b,c,d)&=\frac{(a-ix)_n(b-ix)_n(c-ix)_n(d-ix)_n}{(-2ix)_n}\\ &\times {}_7F_6\left(\begin{matrix}2ix-n,ix-\frac12n+1,a+ix,b+ix,c+ix,d+ix,-n\\ ix-\frac12n,1-n-a+ix,1-n-b+ix,1-n-c+ix,1-n-d+ix\end{matrix};1\right). \end{aligned}\tag{10}$$

The symmetry in a,b,c,d is clear from (10).

Special value

$$ W_n(-a^2;a,b,c,d)=(a+b)_n(a+c)_n(a+d)_n\,, \tag{11}$$

and similarly for arguments $-b^2$, $-c^2$ and $-d^2$ by symmetry of W_n in a,b,c,d.

Uniqueness of orthogonality measure Under the assumptions on a, b, c, d for (9.1.2) or (9.1.3) the orthogonality measure is unique up to constant factor.

For the proof assume without loss of generality (by the symmetry in a, b, c, d) that $\operatorname{Re} a \geq 0$. Write the right-hand side of (9.1.2) or (9.1.3) as $h_n \delta_{m,n}$. Observe from (9.1.2) and (11) that

$$\frac{|W_n(-a^2; a, b, c, d)|^2}{h_n} = O(n^{4\operatorname{Re} a - 1}) \quad \text{as } n \to \infty.$$

Therefore (1) holds, from which the uniqueness of the orthogonality measure follows.

By a similar, but necessarily more complicated argument Ismail et al. [281, Section 3] proved the uniqueness of orthogonality measure for associated Wilson polynomials.

9.2 Racah

Racah in terms of Wilson In the Remark on p.196 Racah polynomials are expressed in terms of Wilson polynomials. This can be equivalently written as

$$\begin{aligned} &R_n\big(x(x-N+\delta); \alpha, \beta, -N-1, \delta\big) \\ &= \frac{W_n\big(-(x+\frac{1}{2}(\delta-N))^2; \frac{1}{2}(\delta-N), \alpha+1-\frac{1}{2}(\delta-N), \beta+\frac{1}{2}(\delta+N)+1, -\frac{1}{2}(\delta+N)\big)}{(\alpha+1)_n(\beta+\delta+1)_n(-N)_n}. \end{aligned} \tag{12}$$

9.3 Continuous dual Hahn

Symmetry The continuous dual Hahn polynomial $S_n(y; a, b, c)$ is symmetric in a, b, c.
This follows from the orthogonality relation (9.3.2) together with the value of its coefficient of y^n given in (9.3.5b). Alternatively, combine (9.3.1) with [AAR, Corollary 3.3.5]].
As a consequence, it is sufficient to give generating function (9.3.12). Then the generating functions (9.3.13), (9.3.14) will follow by symmetry in the parameters.

Special value

$$S_n(-a^2; a, b, c) = (a+b)_n(a+c)_n\,, \tag{13}$$

and similarly for arguments $-b^2$ and $-c^2$ by symmetry of S_n in a, b, c.

Uniqueness of orthogonality measure Under the assumptions on a, b, c for (9.3.2) or (9.3.3) the orthogonality measure is unique up to constant factor.

For the proof assume without loss of generality (by the symmetry in a, b, c, d) that $\operatorname{Re} a \geq 0$. Write the right-hand side of (9.3.2) or (9.3.3) as $h_n \delta_{m,n}$. Observe from (9.3.2) and (13) that

$$\frac{|S_n(-a^2; a, b, c)|^2}{h_n} = O(n^{2\operatorname{Re} a - 1}) \quad \text{as } n \to \infty.$$

Therefore (1) holds, from which the uniqueness of the orthogonality measure follows.

9.4 Continuous Hahn

Orthogonality relation and symmetry The orthogonality relation (9.4.2) holds under the more general assumption that $\mathrm{Re}\,(a,b,c,d) > 0$ and $(c,d) = (\overline{a},\overline{b})$ or $(\overline{b},\overline{a})$.
Thus, under these assumptions, the continuous Hahn polynomial $p_n(x;a,b,c,d)$ is symmetric in a,b and in c,d. This follows from the orthogonality relation (9.4.2) together with the value of its coefficient of x^n given in (9.4.4b).
As a consequence, it is sufficient to give generating function (9.4.11). Then the generating function (9.4.12) will follow by symmetry in the parameters.

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.4.4) behaves as $O(n^2)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

9.5 Hahn

Special values

$$Q_n(0;\alpha,\beta,N) = 1, \quad Q_n(N;\alpha,\beta,N) = \frac{(-1)^n(\beta+1)_n}{(\alpha+1)_n}\,. \tag{14}$$

Use (9.5.1) and compare with (9.8.1) and (33).

From (9.5.3) and (3) it follows that

$$Q_{2n}(N;\alpha,\alpha,2N) = \frac{(\frac{1}{2})_n(N+\alpha+1)_n}{(-N+\frac{1}{2})_n(\alpha+1)_n}\,. \tag{15}$$

From (9.5.1) and [DLMF, (15.4.24)]] it follows that

$$Q_N(x;\alpha,\beta,N) = \frac{(-N-\beta)_x}{(\alpha+1)_x} \qquad (x = 0,1,\ldots,N). \tag{16}$$

Symmetries By the orthogonality relation (9.5.2):

$$\frac{Q_n(N-x;\alpha,\beta,N)}{Q_n(N;\alpha,\beta,N)} = Q_n(x;\beta,\alpha,N), \tag{17}$$

It follows from (24) and (19) that

$$\frac{Q_{N-n}(x;\alpha,\beta,N)}{Q_N(x;\alpha,\beta,N)} = Q_n(x;-N-\beta-1,-N-\alpha-1,N) \qquad (x = 0,1,\ldots,N). \tag{18}$$

Duality The Remark on p.208 gives the duality between Hahn and dual Hahn polynomials:

$$Q_n(x;\alpha,\beta,N) = R_x(n(n+\alpha+\beta+1);\alpha,\beta,N) \quad (n,x \in \{0,1,\ldots N\}). \tag{19}$$

9.6 Dual Hahn

Special values By (16) and (19) we have

$$R_n(N(N+\gamma+\delta+1);\gamma,\delta,N) = \frac{(-N-\delta)_n}{(\gamma+1)_n}\,. \tag{20}$$

It follows from (14) and (19) that

$$R_N(x(x+\gamma+\delta+1);\gamma,\delta,N) = \frac{(-1)^x(\delta+1)_x}{(\gamma+1)_x} \qquad (x=0,1,\ldots,N). \tag{21}$$

Symmetries Write the weight in (9.6.2) as

$$w_x(\alpha,\beta,N) := N!\,\frac{2x+\gamma+\delta+1}{(x+\gamma+\delta+1)_{N+1}}\,\frac{(\gamma+1)_x}{(\delta+1)_x}\binom{N}{x}. \tag{22}$$

Then

$$(\delta+1)_N\,w_{N-x}(\gamma,\delta,N) = (-\gamma-N)_N\,w_x(-\delta-N-1,-\gamma-N-1,N). \tag{23}$$

Hence, by (9.6.2),

$$\frac{R_n((N-x)(N-x+\gamma+\delta+1);\gamma,\delta,N)}{R_n(N(N+\gamma+\delta+1);\gamma,\delta,N)} = R_n(x(x-2N-\gamma-\delta-1);-N-\delta-1,-N-\gamma-1,N). \tag{24}$$

Alternatively, (24) follows from (9.6.1) and [DLMF, (16.4.11)]].

It follows from (17) and (19) that

$$\frac{R_{N-n}(x(x+\gamma+\delta+1);\gamma,\delta,N)}{R_N(x(x+\gamma+\delta+1);\gamma,\delta,N)} = R_n(x(x+\gamma+\delta+1);\delta,\gamma,N) \qquad (x=0,1,\ldots,N). \tag{25}$$

Re: (9.6.11). The generating function (9.6.11) can be written in a more conceptual way as

$$(1-t)^x\;{}_2F_1\left(\begin{matrix}x-N,x+\gamma+1\\-\delta-N\end{matrix};t\right) = \frac{N!}{(\delta+1)_N}\sum_{n=0}^{N}\omega_n\,R_n(\lambda(x);\gamma,\delta,N)\,t^n, \tag{26}$$

where

$$\omega_n := \binom{\gamma+n}{n}\binom{\delta+N-n}{N-n}, \tag{27}$$

i.e., the denominator on the right-hand side of (9.6.2). By the duality between Hahn polynomials and dual Hahn polynomials (see (19)) the above generating function can be rewritten in terms of Hahn polynomials:

$$(1-t)^n\;{}_2F_1\left(\begin{matrix}n-N,n+\alpha+1\\-\beta-N\end{matrix};t\right) = \frac{N!}{(\beta+1)_N}\sum_{x=0}^{N}w_x\,Q_n(x;\alpha,\beta,N)\,t^x, \tag{28}$$

where

$$w_x := \binom{\alpha+x}{x}\binom{\beta+N-x}{N-x}, \tag{29}$$

i.e., the weight occurring in the orthogonality relation (9.5.2) for Hahn polynomials.

Re: (9.6.15). There should be a closing bracket before the equality sign.

9.7 Meixner-Pollaczek

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.7.4) behaves as $O(n^2)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

9.8 Jacobi

Orthogonality relation Write the right-hand side of (9.8.2) as $h_n\,\delta_{m,n}$. Then

$$\begin{aligned}
&\frac{h_n}{h_0} = \frac{n+\alpha+\beta+1}{2n+\alpha+\beta+1}\,\frac{(\alpha+1)_n(\beta+1)_n}{(\alpha+\beta+2)_n\,n!}\,, \quad h_0 = \frac{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(\beta+1)}{\Gamma(\alpha+\beta+2)}\,,\\
&\frac{h_n}{h_0\,(P_n^{(\alpha,\beta)}(1))^2} = \frac{n+\alpha+\beta+1}{2n+\alpha+\beta+1}\,\frac{(\beta+1)_n\,n!}{(\alpha+1)_n\,(\alpha+\beta+2)_n}\,.
\end{aligned} \tag{30}$$

In (9.8.3) the numerator factor $\Gamma(n+\alpha+\beta+1)$ in the last line should be $\Gamma(\beta+1)$. When thus corrected, (9.8.3) can be rewritten as:

$$\begin{aligned}
&\int_1^\infty P_m^{(\alpha,\beta)}(x)\,P_n^{(\alpha,\beta)}(x)\,(x-1)^\alpha(x+1)^\beta\,dx = h_n\,\delta_{m,n}\,,\\
&\qquad\qquad -1-\beta > \alpha > -1, \quad m,n < -\tfrac{1}{2}(\alpha+\beta+1),\\
&\frac{h_n}{h_0} = \frac{n+\alpha+\beta+1}{2n+\alpha+\beta+1}\,\frac{(\alpha+1)_n(\beta+1)_n}{(\alpha+\beta+2)_n\,n!}\,, \quad h_0 = \frac{2^{\alpha+\beta+1}\Gamma(\alpha+1)\Gamma(-\alpha-\beta-1)}{\Gamma(-\beta)}\,.
\end{aligned} \tag{31}$$

Symmetry

$$P_n^{(\alpha,\beta)}(-x) = (-1)^n\,P_n^{(\beta,\alpha)}(x). \tag{32}$$

Use (9.8.2) and (9.8.5b) or see [DLMF, Table 18.6.1]].

Special values

$$P_n^{(\alpha,\beta)}(1) = \frac{(\alpha+1)_n}{n!}\,, \quad P_n^{(\alpha,\beta)}(-1) = \frac{(-1)^n(\beta+1)_n}{n!}\,, \quad \frac{P_n^{(\alpha,\beta)}(-1)}{P_n^{(\alpha,\beta)}(1)} = \frac{(-1)^n(\beta+1)_n}{(\alpha+1)_n}\,. \tag{33}$$

Use (9.8.1) and (32) or see [DLMF, Table 18.6.1]].

Generating functions Formula (9.8.15) was first obtained by Brafman [109].

Bilateral generating functions For $0 \le r < 1$ and $x, y \in [-1, 1]$ we have in terms of F_4 (see (4)):

$$\sum_{n=0}^{\infty} \frac{(\alpha+\beta+1)_n\, n!}{(\alpha+1)_n(\beta+1)_n}\, r^n\, P_n^{(\alpha,\beta)}(x)\, P_n^{(\alpha,\beta)}(y) = \frac{1}{(1+r)^{\alpha+\beta+1}} \times F_4\Big(\tfrac{1}{2}(\alpha+\beta+1), \tfrac{1}{2}(\alpha+\beta+2); \alpha+1, \beta+1; \frac{r(1-x)(1-y)}{(1+r)^2}, \frac{r(1+x)(1+y)}{(1+r)^2}\Big), \quad (34)$$

$$\sum_{n=0}^{\infty} \frac{2n+\alpha+\beta+1}{n+\alpha+\beta+1} \frac{(\alpha+\beta+2)_n\, n!}{(\alpha+1)_n(\beta+1)_n}\, r^n\, P_n^{(\alpha,\beta)}(x)\, P_n^{(\alpha,\beta)}(y) = \frac{1-r}{(1+r)^{\alpha+\beta+2}} \times F_4\Big(\tfrac{1}{2}(\alpha+\beta+2), \tfrac{1}{2}(\alpha+\beta+3); \alpha+1, \beta+1; \frac{r(1-x)(1-y)}{(1+r)^2}, \frac{r(1+x)(1+y)}{(1+r)^2}\Big). \quad (35)$$

Formulas (34) and (35) were first given by Bailey [91, (2.1), (2.3)]. See Stanton [485] for a shorter proof. (However, in the second line of [485, (1)] z and Z should be interchanged.) As observed in Bailey [91, p.10], (35) follows from (34) by applying the operator $r^{-\frac{1}{2}(\alpha+\beta-1)}\, \frac{d}{dr} \circ r^{\frac{1}{2}(\alpha+\beta+1)}$ to both sides of (34). In view of (30), formula (35) is the Poisson kernel for Jacobi polynomials. The right-hand side of (35) makes clear that this kernel is positive. See also the discussion in Askey [46, following (2.32)].

Quadratic transformations

$$\frac{C_{2n}^{(\alpha+\frac{1}{2})}(x)}{C_{2n}^{(\alpha+\frac{1}{2})}(1)} = \frac{P_{2n}^{(\alpha,\alpha)}(x)}{P_{2n}^{(\alpha,\alpha)}(1)} = \frac{P_n^{(\alpha,-\frac{1}{2})}(2x^2-1)}{P_n^{(\alpha,-\frac{1}{2})}(1)}, \quad (36)$$

$$\frac{C_{2n+1}^{(\alpha+\frac{1}{2})}(x)}{C_{2n+1}^{(\alpha+\frac{1}{2})}(1)} = \frac{P_{2n+1}^{(\alpha,\alpha)}(x)}{P_{2n+1}^{(\alpha,\alpha)}(1)} = \frac{x\, P_n^{(\alpha,\frac{1}{2})}(2x^2-1)}{P_n^{(\alpha,\frac{1}{2})}(1)}. \quad (37)$$

See p.221, Remarks, last two formulas together with (33) and (48). Or see [DLMF, (18.7.13), (18.7.14)]].

Differentiation formulas Each differentiation formula is given in two equivalent forms.

$$\begin{aligned} \frac{d}{dx}\Big((1-x)^{\alpha} P_n^{(\alpha,\beta)}(x)\Big) &= -(n+\alpha)\,(1-x)^{\alpha-1} P_n^{(\alpha-1,\beta+1)}(x), \\ \Big((1-x)\frac{d}{dx} - \alpha\Big)\, P_n^{(\alpha,\beta)}(x) &= -(n+\alpha)\, P_n^{(\alpha-1,\beta+1)}(x). \end{aligned} \quad (38)$$

$$\begin{aligned} \frac{d}{dx}\Big((1+x)^{\beta} P_n^{(\alpha,\beta)}(x)\Big) &= (n+\beta)\,(1+x)^{\beta-1} P_n^{(\alpha+1,\beta-1)}(x), \\ \Big((1+x)\frac{d}{dx} + \beta\Big)\, P_n^{(\alpha,\beta)}(x) &= (n+\beta)\, P_n^{(\alpha+1,\beta-1)}(x). \end{aligned} \quad (39)$$

Formulas (38) and (39) follow from [DLMF, (15.5.4), (15.5.6)]] together with (9.8.1). They also follow from each other by (32).

Generalized Gegenbauer polynomials See [146, p.156]. These are defined by

$$S_{2m}^{(\alpha,\beta)}(x) := \text{const.}\, P_m^{(\alpha,\beta)}(2x^2-1), \qquad S_{2m+1}^{(\alpha,\beta)}(x) := \text{const.}\, x\, P_m^{(\alpha,\beta+1)}(2x^2-1). \tag{40}$$

Then for $\alpha, \beta > -1$ we have the orthogonality relation

$$\int_{-1}^{1} S_m^{(\alpha,\beta)}(x)\, S_n^{(\alpha,\beta)}(x)\, |x|^{2\beta+1}(1-x^2)^\alpha\, dx = 0 \qquad (m \neq n). \tag{41}$$

If we define the *Dunkl operator* T_μ by

$$(T_\mu f)(x) := f'(x) + \mu\, \frac{f(x)-f(-x)}{x} \tag{42}$$

and if we choose the constants in (40) as

$$S_{2m}^{(\alpha,\beta)}(x) = \frac{(\alpha+\beta+1)_m}{(\beta+1)_m}\, P_m^{(\alpha,\beta)}(2x^2-1), \quad S_{2m+1}^{(\alpha,\beta)}(x) = \frac{(\alpha+\beta+1)_{m+1}}{(\beta+1)_{m+1}}\, x\, P_m^{(\alpha,\beta+1)}(2x^2-1) \tag{43}$$

then (see [K5, (1.6)])

$$T_{\beta+\frac{1}{2}} S_n^{(\alpha,\beta)} = 2(\alpha+\beta+1)\, S_{n-1}^{(\alpha+1,\beta)}. \tag{44}$$

Formula (44) with (43) substituted gives rise to two differentiation formulas involving Jacobi polynomials which are equivalent to (9.8.7) and (39).

Composition of (44) with itself gives

$$T_{\beta+\frac{1}{2}}^2 S_n^{(\alpha,\beta)} = 4(\alpha+\beta+1)(\alpha+\beta+2)\, S_{n-2}^{(\alpha+2,\beta)},$$

which is equivalent to the composition of (9.8.7) and (39):

$$\left(\frac{d^2}{dx^2} + \frac{2\beta+1}{x}\,\frac{d}{dx}\right) P_n^{(\alpha,\beta)}(2x^2-1) = 4(n+\alpha+\beta+1)(n+\beta)\, P_{n-1}^{(\alpha+2,\beta)}(2x^2-1). \tag{45}$$

Formula (45) was also given in [322, (2.4)].

9.8.1 Gegenbauer / Ultraspherical

Notation Here the Gegenbauer polynomial is denoted by C_n^λ instead of $C_n^{(\lambda)}$.

Orthogonality relation Write the right-hand side of (9.8.20) as $h_n\, \delta_{m,n}$. Then

$$\frac{h_n}{h_0} = \frac{\lambda}{\lambda+n}\,\frac{(2\lambda)_n}{n!}, \quad h_0 = \frac{\pi^{\frac{1}{2}}\,\Gamma(\lambda+\frac{1}{2})}{\Gamma(\lambda+1)}, \quad \frac{h_n}{h_0\,(C_n^\lambda(1))^2} = \frac{\lambda}{\lambda+n}\,\frac{n!}{(2\lambda)_n}. \tag{46}$$

Hypergeometric representation Beside (9.8.19) we have also

$$C_n^\lambda(x) = \sum_{\ell=0}^{\lfloor n/2\rfloor} \frac{(-1)^\ell(\lambda)_{n-\ell}}{\ell!\,(n-2\ell)!}\,(2x)^{n-2\ell} = (2x)^n\,\frac{(\lambda)_n}{n!}\;{}_2F_1\left(\begin{matrix}-\frac{1}{2}n, -\frac{1}{2}n+\frac{1}{2}\\ 1-\lambda-n\end{matrix}; \frac{1}{x^2}\right). \tag{47}$$

See [DLMF, (18.5.10)]].

Special value

$$C_n^\lambda(1) = \frac{(2\lambda)_n}{n!}\,. \tag{48}$$

Use (9.8.19) or see [DLMF, Table 18.6.1]].

Expression in terms of Jacobi

$$\frac{C_n^\lambda(x)}{C_n^\lambda(1)} = \frac{P_n^{(\lambda-\frac12,\lambda-\frac12)}(x)}{P_n^{(\lambda-\frac12,\lambda-\frac12)}(1)}\,, \qquad C_n^\lambda(x) = \frac{(2\lambda)_n}{(\lambda+\frac12)_n}\,P_n^{(\lambda-\frac12,\lambda-\frac12)}(x). \tag{49}$$

Re: (9.8.21) By iteration of recurrence relation (9.8.21):

$$x^2C_n^\lambda(x) = \frac{(n+1)(n+2)}{4(n+\lambda)(n+\lambda+1)}\,C_{n+2}^\lambda(x) + \frac{n^2+2n\lambda+\lambda-1}{2(n+\lambda-1)(n+\lambda+1)}\,C_n^\lambda(x) + \frac{(n+2\lambda-1)(n+2\lambda-2)}{4(n+\lambda)(n+\lambda-1)}\,C_{n-2}^\lambda(x). \tag{50}$$

Bilateral generating functions

$$\sum_{n=0}^\infty \frac{n!}{(2\lambda)_n}\,r^n\,C_n^\lambda(x)\,C_n^\lambda(y) = \frac{1}{(1-2rxy+r^2)^\lambda}\;{}_2F_1\left(\begin{matrix}\frac12\lambda,\frac12(\lambda+1)\\ \lambda+\frac12\end{matrix};\frac{4r^2(1-x^2)(1-y^2)}{(1-2rxy+r^2)^2}\right) \qquad (r\in(-1,1),\ x,y\in[-1,1]). \tag{51}$$

For the proof put $\beta := \alpha$ in (34), then use (5) and (49). The Poisson kernel for Gegenbauer polynomials can be derived in a similar way from (35), or alternatively by applying the operator $r^{-\lambda+1}\frac{d}{dr}\circ r^\lambda$ to both sides of (51):

$$\sum_{n=0}^\infty \frac{\lambda+n}{\lambda}\,\frac{n!}{(2\lambda)_n}\,r^n\,C_n^\lambda(x)\,C_n^\lambda(y) = \frac{1-r^2}{(1-2rxy+r^2)^{\lambda+1}} \times {}_2F_1\left(\begin{matrix}\frac12(\lambda+1),\frac12(\lambda+2)\\ \lambda+\frac12\end{matrix};\frac{4r^2(1-x^2)(1-y^2)}{(1-2rxy+r^2)^2}\right) \qquad (r\in(-1,1),\ x,y\in[-1,1]). \tag{52}$$

Formula (52) was obtained by Gasper & Rahman [234, (4.4)] as a limit case of their formula for the Poisson kernel for continuous q-ultraspherical polynomials.

Trigonometric expansions By [DLMF, (18.5.11), (15.8.1)]]:

$$C_n^\lambda(\cos\theta) = \sum_{k=0}^n \frac{(\lambda)_k(\lambda)_{n-k}}{k!\,(n-k)!}\,e^{i(n-2k)\theta} = e^{in\theta}\frac{(\lambda)_n}{n!}\;{}_2F_1\left(\begin{matrix}-n,\lambda\\ 1-\lambda-n\end{matrix};e^{-2i\theta}\right) \tag{53}$$

$$= \frac{(\lambda)_n}{2^\lambda n!}\,e^{-\frac12 i\lambda\pi}e^{i(n+\lambda)\theta}\,(\sin\theta)^{-\lambda}\;{}_2F_1\left(\begin{matrix}\lambda,1-\lambda\\ 1-\lambda-n\end{matrix};\frac{ie^{-i\theta}}{2\sin\theta}\right) \tag{54}$$

$$= \frac{(\lambda)_n}{n!}\sum_{k=0}^\infty \frac{(\lambda)_k(1-\lambda)_k}{(1-\lambda-n)_k k!}\,\frac{\cos((n-k+\lambda)\theta+\frac12(k-\lambda)\pi)}{(2\sin\theta)^{k+\lambda}}\,. \tag{55}$$

In (54) and (55) we require that $\frac{1}{6}\pi < \theta < \frac{5}{6}\pi$. Then the convergence is absolute for $\lambda > \frac{1}{2}$ and conditional for $0 < \lambda \le \frac{1}{2}$.

By [DLMF, (14.13.1), (14.3.21), (15.8.1)]]:

$$C_n^\lambda(\cos\theta) = \frac{2\Gamma(\lambda+\frac{1}{2})}{\pi^{\frac{1}{2}}\Gamma(\lambda+1)}\,\frac{(2\lambda)_n}{(\lambda+1)_n}\,(\sin\theta)^{1-2\lambda}\sum_{k=0}^{\infty}\frac{(1-\lambda)_k(n+1)_k}{(n+\lambda+1)_k k!}\,\sin\big((2k+n+1)\theta\big) \tag{56}$$

$$= \frac{2\Gamma(\lambda+\frac{1}{2})}{\pi^{\frac{1}{2}}\Gamma(\lambda+1)}\,\frac{(2\lambda)_n}{(\lambda+1)_n}\,(\sin\theta)^{1-2\lambda}\,\mathrm{Im}\left(e^{i(n+1)\theta}\;{}_2F_1\left(\begin{matrix}1-\lambda, n+1\\ n+\lambda+1\end{matrix}; e^{2i\theta}\right)\right)$$

$$= \frac{2^\lambda\Gamma(\lambda+\frac{1}{2})}{\pi^{\frac{1}{2}}\Gamma(\lambda+1)}\,\frac{(2\lambda)_n}{(\lambda+1)_n}\,(\sin\theta)^{-\lambda}\,\mathrm{Re}\left(e^{-\frac{1}{2}i\lambda\pi}e^{i(n+\lambda)\theta}\;{}_2F_1\left(\begin{matrix}\lambda, 1-\lambda\\ 1+\lambda+n\end{matrix}; \frac{e^{i\theta}}{2i\sin\theta}\right)\right)$$

$$= \frac{2^{2\lambda}\Gamma(\lambda+\frac{1}{2})}{\pi^{\frac{1}{2}}\Gamma(\lambda+1)}\,\frac{(2\lambda)_n}{(\lambda+1)_n}\sum_{k=0}^{\infty}\frac{(\lambda)_k(1-\lambda)_k}{(1+\lambda+n)_k k!}\,\frac{\cos((n+k+\lambda)\theta-\frac{1}{2}(k+\lambda)\pi)}{(2\sin\theta)^{k+\lambda}}\,. \tag{57}$$

We require that $0 < \theta < \pi$ in (56) and $\frac{1}{6}\pi < \theta < \frac{5}{6}\pi$ in (57) The convergence is absolute for $\lambda > \frac{1}{2}$ and conditional for $0 < \lambda \le \frac{1}{2}$. For $\lambda \in \mathbb{Z}_{>0}$ the above series terminate after the term with $k = \lambda - 1$. Formulas (56) and (57) are also given in [Sz, (4.9.22), (4.9.25)].

Fourier transform

$$\frac{\Gamma(\lambda+1)}{\Gamma(\lambda+\frac{1}{2})\,\Gamma(\frac{1}{2})}\int_{-1}^{1}\frac{C_n^\lambda(y)}{C_n^\lambda(1)}\,(1-y^2)^{\lambda-\frac{1}{2}}\,e^{ixy}\,dy = i^n\,2^\lambda\,\Gamma(\lambda+1)\,x^{-\lambda}\,J_{\lambda+n}(x). \tag{58}$$

See [DLMF, (18.17.17) and (18.17.18)]].

Laplace transforms

$$\frac{2}{n!\,\Gamma(\lambda)}\int_0^\infty H_n(tx)\,t^{n+2\lambda-1}\,e^{-t^2}\,dt = C_n^\lambda(x). \tag{59}$$

See Nielsen [K20, p.48, (4) with p.47, (1) and p.28, (10)] (1918) or Feldheim [K8, (28)] (1942).

$$\frac{2}{\Gamma(\lambda+\frac{1}{2})}\int_0^1\frac{C_n^\lambda(t)}{C_n^\lambda(1)}\,(1-t^2)^{\lambda-\frac{1}{2}}\,t^{-1}\,(x/t)^{n+2\lambda+1}\,e^{-x^2/t^2}\,dt = 2^{-n}\,H_n(x)\,e^{-x^2}\quad(\lambda > -\tfrac{1}{2}). \tag{60}$$

Use Askey & Fitch [K1, (3.29)] for $\alpha = \pm\frac{1}{2}$ together with (32), (36), (37), (82) and (83).

Addition formula (see [AAR, (9.8.5′)]])

$$R_n^{(\alpha,\alpha)}\big(xy+(1-x^2)^{\frac{1}{2}}(1-y^2)^{\frac{1}{2}}t\big) = \sum_{k=0}^{n}\frac{(-1)^k(-n)_k\,(n+2\alpha+1)_k}{2^{2k}((\alpha+1)_k)^2}$$

$$\times\,(1-x^2)^{k/2}R_{n-k}^{(\alpha+k,\alpha+k)}(x)\,(1-y^2)^{k/2}R_{n-k}^{(\alpha+k,\alpha+k)}(y)\,\omega_k^{(\alpha-\frac{1}{2},\alpha-\frac{1}{2})}\,R_k^{(\alpha-\frac{1}{2},\alpha-\frac{1}{2})}(t), \tag{61}$$

where

$$R_n^{(\alpha,\beta)}(x) := P_n^{(\alpha,\beta)}(x)/P_n^{(\alpha,\beta)}(1),\quad \omega_n^{(\alpha,\beta)} := \frac{\int_{-1}^1(1-x)^\alpha(1+x)^\beta\,dx}{\int_{-1}^1(R_n^{(\alpha,\beta)}(x))^2\,(1-x)^\alpha(1+x)^\beta\,dx}\,.$$

9.8.2 Chebyshev

In addition to the Chebyshev polynomials T_n of the first kind (9.8.35) and U_n of the second kind (9.8.36),

$$T_n(x) := \frac{P_n^{(-\frac{1}{2},-\frac{1}{2})}(x)}{P_n^{(-\frac{1}{2},-\frac{1}{2})}(1)} = \cos(n\theta), \quad x = \cos\theta, \tag{62}$$

$$U_n(x) := (n+1)\,\frac{P_n^{(\frac{1}{2},\frac{1}{2})}(x)}{P_n^{(\frac{1}{2},\frac{1}{2})}(1)} = \frac{\sin((n+1)\theta)}{\sin\theta}, \quad x = \cos\theta, \tag{63}$$

we have Chebyshev polynomials V_n *of the third kind* and W_n *of the fourth kind*,

$$V_n(x) := \frac{P_n^{(-\frac{1}{2},\frac{1}{2})}(x)}{P_n^{(-\frac{1}{2},\frac{1}{2})}(1)} = \frac{\cos((n+\frac{1}{2})\theta)}{\cos(\frac{1}{2}\theta)}, \quad x = \cos\theta, \tag{64}$$

$$W_n(x) := (2n+1)\,\frac{P_n^{(\frac{1}{2},-\frac{1}{2})}(x)}{P_n^{(\frac{1}{2},-\frac{1}{2})}(1)} = \frac{\sin((n+\frac{1}{2})\theta)}{\sin(\frac{1}{2}\theta)}, \quad x = \cos\theta, \tag{65}$$

see [K18, Section 1.2.3]. Then there is the symmetry

$$V_n(-x) = (-1)^n W_n(x). \tag{66}$$

The names of Chebyshev polynomials of the third and fourth kind and the notation $V_n(x)$ are due to Gautschi [K9]. The notation $W_n(x)$ was first used by Mason [K17]. Names and notations for Chebyshev polynomials of the third and fourth kind are interchanged in [AAR, Remark 2.5.3]] and [DLMF, Table 18.3.1]].

9.9 Pseudo Jacobi (or Routh-Romanovski)

In this section in the book the pseudo Jacobi polynomial $P_n(x;\nu,N)$ in (9.9.1) is considered for $N \in \mathbb{Z}_{\geq 0}$ and $n = 0, 1, \ldots, n$. However, we can more generally take $-\frac{1}{2} < N \in \mathbb{R}$ (so here I overrule my convention formulated in the beginning of this paper), N_0 integer such that $N - \frac{1}{2} \leq N_0 < N + \frac{1}{2}$, and $n = 0, 1, \ldots, N_0$ (see [382, §5, case A.4]). The orthogonality relation (9.9.2) is valid for $m, n = 0, 1, \ldots, N_0$.

History These polynomials were first obtained by Routh [K23] in 1885, and later, independently, by Romanovski [463] in 1929.

Limit relation: Pseudo big q-Jacobi $\longrightarrow$ Pseudo Jacobi
See also (131).

References See also [Ism, §20.1], [51], [384], [K13], [K16], [K21].

9.10 Meixner

History In 1934 Meixner [406] (see (1.1) and case IV on pp. 10, 11 and 12) gave the orthogonality measure for the polynomials P_n given by the generating function

$$e^{xu(t)}\, f(t) = \sum_{n=0}^{\infty} P_n(x)\, \frac{t^n}{n!}\,,$$

where

$$e^{u(t)} = \left(\frac{1-\beta t}{1-\alpha t}\right)^{\frac{1}{\alpha-\beta}}, \quad f(t) = \frac{(1-\beta t)^{\frac{k_2}{\beta(\alpha-\beta)}}}{(1-\alpha t)^{\frac{k_2}{\alpha(\alpha-\beta)}}} \quad (k_2 < 0;\ \alpha > \beta > 0 \ \text{ or } \ \alpha < \beta < 0).$$

Then P_n can be expressed as a Meixner polynomial:

$$P_n(x) = (-k_2(\alpha\beta)^{-1})_n\, \beta^n\, M_n\left(-\frac{x + k_2\alpha^{-1}}{\alpha-\beta}, -k_2(\alpha\beta)^{-1}, \beta\alpha^{-1}\right).$$

In 1938 Gottlieb [K11, §2] introduces polynomials l_n "of Laguerre type" which turn out to be special Meixner polynomials: $l_n(x) = e^{-n\lambda} M_n(x; 1, e^{-\lambda})$.

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.10.4) behaves as $O(n^2)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

9.11 Krawtchouk

Special values By (9.11.1) and the binomial formula:

$$K_n(0; p, N) = 1, \qquad K_n(N; p, N) = (1 - p^{-1})^n. \tag{67}$$

The self-duality (p.240, Remarks, first formula)

$$K_n(x; p, N) = K_x(n; p, N) \qquad (n, x \in \{0, 1, \ldots, N\}) \tag{68}$$

combined with (67) yields:

$$K_N(x; p, N) = (1 - p^{-1})^x \qquad (x \in \{0, 1, \ldots, N\}). \tag{69}$$

Symmetry By the orthogonality relation (9.11.2):

$$\frac{K_n(N - x; p, N)}{K_n(N; p, N)} = K_n(x; 1 - p, N). \tag{70}$$

By (70) and (68) we have also

$$\frac{K_{N-n}(x; p, N)}{K_N(x; p, N)} = K_n(x; 1 - p, N) \qquad (n, x \in \{0, 1, \ldots, N\}), \tag{71}$$

and, by (71), (70) and (67),

$$K_{N-n}(N-x;p,N)=\left(\frac{p}{p-1}\right)^{n+x-N}K_n(x;p.N)\qquad(n,x\in\{0,1,\ldots,N\}).\tag{72}$$

A particular case of (70) is:

$$K_n(N-x;\tfrac{1}{2},N)=(-1)^nK_n(x;\tfrac{1}{2},N).\tag{73}$$

Hence

$$K_{2m+1}(N;\tfrac{1}{2},2N)=0.\tag{74}$$

From (9.11.11):

$$K_{2m}(N;\tfrac{1}{2},2N)=\frac{(\frac{1}{2})_m}{(-N+\frac{1}{2})_m}\,.\tag{75}$$

Quadratic transformations

$$K_{2m}(x+N;\tfrac{1}{2},2N)=\frac{(\frac{1}{2})_m}{(-N+\frac{1}{2})_m}\,R_m(x^2;-\tfrac{1}{2},-\tfrac{1}{2},N),\tag{76}$$

$$K_{2m+1}(x+N;\tfrac{1}{2},2N)=-\frac{(\frac{3}{2})_m}{N\,(-N+\frac{1}{2})_m}\,x\,R_m(x^2-1;\tfrac{1}{2},\tfrac{1}{2},N-1),\tag{77}$$

$$K_{2m}(x+N+1;\tfrac{1}{2},2N+1)=\frac{(\frac{1}{2})_m}{(-N-\frac{1}{2})_m}\,R_m(x(x+1);-\tfrac{1}{2},\tfrac{1}{2},N),\tag{78}$$

$$K_{2m+1}(x+N+1;\tfrac{1}{2},2N+1)=\frac{(\frac{3}{2})_m}{(-N-\frac{1}{2})_{m+1}}\,(x+\tfrac{1}{2})\,R_m(x(x+1);\tfrac{1}{2},-\tfrac{1}{2},N),\tag{79}$$

where R_m is a dual Hahn polynomial (9.6.1). For the proofs use (9.6.2), (9.11.2), (9.6.4) and (9.11.4).

Generating functions

$$\begin{aligned}&\sum_{x=0}^{N}\binom{N}{x}K_m(x;p,N)K_n(x;q,N)z^x\\&\quad=\left(\frac{p-z+pz}{p}\right)^m\left(\frac{q-z+qz}{q}\right)^n(1+z)^{N-m-n}K_m\left(n;-\frac{(p-z+pz)(q-z+qz)}{z},N\right).\end{aligned}\tag{80}$$

This follows immediately from Rosengren [K22, (3.5)], which goes back to Meixner [K19].

9.12 Laguerre

Notation Here the Laguerre polynomial is denoted by L_n^α instead of $L_n^{(\alpha)}$.

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.12.4) behaves as $O(n^2)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

Special value

$$L_n^\alpha(0) = \frac{(\alpha+1)_n}{n!}\,. \tag{81}$$

Use (9.12.1) or see [DLMF, 18.6.1)]].

Quadratic transformations

$$H_{2n}(x) = (-1)^n\, 2^{2n}\, n!\, L_n^{-1/2}(x^2), \tag{82}$$

$$H_{2n+1}(x) = (-1)^n\, 2^{2n+1}\, n!\, x\, L_n^{1/2}(x^2). \tag{83}$$

See p.244, Remarks, last two formulas. Or see [DLMF, (18.7.19), (18.7.20)]].

Fourier transform

$$\frac{1}{\Gamma(\alpha+1)} \int_0^\infty \frac{L_n^\alpha(y)}{L_n^\alpha(0)}\, e^{-y}\, y^\alpha\, e^{ixy}\, dy = i^n\, \frac{y^n}{(iy+1)^{n+\alpha+1}}\,, \tag{84}$$

see [DLMF, (18.17.34)]].

Differentiation formulas Each differentiation formula is given in two equivalent forms.

$$\frac{d}{dx}\,(x^\alpha L_n^\alpha(x)) = (n+\alpha)\, x^{\alpha-1} L_n^{\alpha-1}(x), \qquad \left(x\frac{d}{dx} + \alpha\right) L_n^\alpha(x) = (n+\alpha)\, L_n^{\alpha-1}(x). \tag{85}$$

$$\frac{d}{dx}\left(e^{-x} L_n^\alpha(x)\right) = -e^{-x} L_n^{\alpha+1}(x), \qquad \left(\frac{d}{dx} - 1\right) L_n^\alpha(x) = -L_n^{\alpha+1}(x). \tag{86}$$

Formulas (85) and (86) follow from [DLMF, (13.3.18), (13.3.20)]] together with (9.12.1).

Generalized Hermite polynomials See [146, p.156]. These are defined by

$$H_{2m}^\mu(x) := \text{const.}\, L_m^{\mu-\frac{1}{2}}(x^2), \qquad H_{2m+1}^\mu(x) := \text{const.}\, x\, L_m^{\mu+\frac{1}{2}}(x^2). \tag{87}$$

Then for $\mu > -\frac{1}{2}$ we have orthogonality relation

$$\int_{-\infty}^{\infty} H_m^\mu(x)\, H_n^\mu(x)\, |x|^{2\mu} e^{-x^2}\, dx = 0 \qquad (m \neq n). \tag{88}$$

Let the Dunkl operator T_μ be defined by (42). If we choose the constants in (87) as

$$H_{2m}^\mu(x) = \frac{(-1)^m (2m)!}{(\mu+\frac{1}{2})_m}\, L_m^{\mu-\frac{1}{2}}(x^2), \qquad H_{2m+1}^\mu(x) = \frac{(-1)^m (2m+1)!}{(\mu+\frac{1}{2})_{m+1}}\, x\, L_m^{\mu+\frac{1}{2}}(x^2) \tag{89}$$

then (see [K5, (1.6)])

$$T_\mu H_n^\mu = 2n\, H_{n-1}^\mu. \tag{90}$$

Formula (90) with (89) substituted gives rise to two differentiation formulas involving Laguerre polynomials which are equivalent to (9.12.6) and (85).

Composition of (90) with itself gives

$$T_\mu^2 H_n^\mu = 4n(n-1)\, H_{n-2}^\mu,$$

which is equivalent to the composition of (9.12.6) and (85):

$$\left(\frac{d^2}{dx^2} + \frac{2\alpha+1}{x}\,\frac{d}{dx}\right) L_n^\alpha(x^2) = -4(n+\alpha)\, L_{n-1}^\alpha(x^2). \tag{91}$$

9.14 Charlier

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.14.4) behaves as $O(n)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

9.15 Hermite

Uniqueness of orthogonality measure The coefficient of $p_{n-1}(x)$ in (9.15.4) behaves as $O(n)$ as $n \to \infty$. Hence (2) holds, by which the orthogonality measure is unique.

Fourier transforms

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{\infty} H_n(y)\, e^{-\frac{1}{2}y^2}\, e^{ixy}\, dy = i^n\, H_n(x)\, e^{-\frac{1}{2}x^2}, \tag{92}$$

see [AAR, (6.1.15) and Exercise 6.11].

$$\frac{1}{\sqrt{\pi}} \int_{-\infty}^{\infty} H_n(y)\, e^{-y^2}\, e^{ixy}\, dy = i^n\, x^n\, e^{-\frac{1}{4}x^2}, \tag{93}$$

see [DLMF, (18.17.35)]].

$$\frac{i^n}{2\sqrt{\pi}} \int_{-\infty}^{\infty} y^n\, e^{-\frac{1}{4}y^2}\, e^{-ixy}\, dy = H_n(x)\, e^{-x^2}, \tag{94}$$

see [AAR, (6.1.4)].

14.1 Askey-Wilson

Symmetry The Askey-Wilson polynomials $p_n(x; a, b, c, d \mid q)$ are symmetric in a, b, c, d.

This follows from the orthogonality relation (14.1.2) together with the value of its coefficient of x^n given in (14.1.5b). Alternatively, combine (14.1.1) with [GR, (III.15)].

As a consequence, it is sufficient to give generating function (14.1.13). Then the generating functions (14.1.14), (14.1.15) will follow by symmetry in the parameters.

Basic hypergeometric representation In addition to (14.1.1) we have (in notation (8)):

$$p_n(\cos\theta; a, b, c, d \mid q) = \frac{(ae^{-i\theta}, be^{-i\theta}, ce^{-i\theta}, de^{-i\theta}; q)_n}{(e^{-2i\theta}; q)_n}\, e^{in\theta} \times {}_8W_7\big(q^{-n}e^{2i\theta}; ae^{i\theta}, be^{i\theta}, ce^{i\theta}, de^{i\theta}, q^{-n}; q, q^{2-n}/(abcd)\big). \quad (95)$$

This follows from (14.1.1) by combining (III.15) and (III.19) in [GR]. It is also given in [513, (4.2)], but be aware for some slight errors. The symmetry in a, b, c, d is evident from (95).

Special value

$$p_n\big(\tfrac{1}{2}(a + a^{-1}); a, b, c, d \mid q\big) = a^{-n}\,(ab, ac, ad; q)_n\,, \quad (96)$$

and similarly for arguments $\frac{1}{2}(b+b^{-1})$, $\frac{1}{2}(c+c^{-1})$ and $\frac{1}{2}(d+d^{-1})$ by symmetry of p_n in a, b, c, d.

Trivial symmetry

$$p_n(-x; a, b, c, d \mid q) = (-1)^n p_n(x; -a, -b, -c, -d \mid q). \quad (97)$$

Both (96) and (97) are obtained from (14.1.1).

Re: (14.1.5) Let

$$p_n(x) := \frac{p_n(x; a, b, c, d \mid q)}{2^n(abcdq^{n-1}; q)_n} = x^n + \widetilde{k}_n x^{n-1} + \cdots\,. \quad (98)$$

Then

$$\widetilde{k}_n = -\frac{(1-q^n)(a+b+c+d-(abc+abd+acd+bcd)q^{n-1})}{2(1-q)(1-abcdq^{2n-2})}\,. \quad (99)$$

This follows because $\tilde{k}_n - \tilde{k}_{n+1}$ equals the coefficient $\frac{1}{2}\big(a+a^{-1}-(A_n+C_n)\big)$ of $p_n(x)$ in (14.1.5).

Limit relations

Askey-Wilson $\longrightarrow$ Wilson
Instead of (14.1.21) we can keep a polynomial of degree n while the limit is approached:

$$\lim_{q\to 1} \frac{p_n(1-\frac{1}{2}x(1-q)^2; q^a, q^b, q^c, q^d \mid q)}{(1-q)^{3n}} = W_n(x; a, b, c, d). \quad (100)$$

For the proof first derive the corresponding limit for the monic polynomials by comparing (14.1.5) with (9.1.5).

References See also Koornwinder [K14].

14.2 q-Racah

Symmetry

$$R_n(x;\alpha,\beta,q^{-N-1},\delta \mid q) = \frac{(\beta q,\alpha\delta^{-1}q;q)_n}{(\alpha q,\beta\delta q;q)_n}\,\delta^n\,R_n(\delta^{-1}x;\beta,\alpha,q^{-N-1},\delta^{-1} \mid q). \tag{101}$$

This follows from (14.2.1) combined with [GR, (III.15)].
In particular,

$$R_n(x;\alpha,\beta,q^{-N-1},-1 \mid q) = \frac{(\beta q,-\alpha q;q)_n}{(\alpha q,-\beta q;q)_n}\,(-1)^n\,R_n(-x;\beta,\alpha,q^{-N-1},-1 \mid q), \tag{102}$$

and

$$R_n(x;\alpha,\alpha,q^{-N-1},-1 \mid q) = (-1)^n\,R_n(-x;\alpha,\alpha,q^{-N-1},-1 \mid q), \tag{103}$$

Trivial symmetry Clearly from (14.2.1):

$$R_n(y;\alpha,\beta,\gamma,\delta \mid q) = R_n(y;\beta\delta,\alpha\delta^{-1},\gamma,\delta \mid q). \tag{104}$$

14.3 Continuous dual q-Hahn

The continuous dual q-Hahn polynomials are the special case $d = 0$ of the Askey-Wilson polynomials:

$$p_n(x;a,b,c \mid q) := p_n(x;a,b,c,0 \mid q).$$

Hence all formulas in §14.3 are specializations for $d = 0$ of formulas in §14.1.

14.4 Continuous q-Hahn

The continuous q-Hahn polynomials are a special case of Askey-Wilson polynomials. Following [72, (4.39)] and [GR, (7.5.43)] we denote these polynomials, instead of $p_n(\cos(\theta+\phi);a,b,a,b;q)$, by

$$p_n(x;a,b \mid q) := p_n(x;ae^{i\phi},be^{i\phi},ae^{-i\phi},be^{-i\phi} \mid q) \quad (x=\cos(\theta+\phi)).$$

(Curiously, this notation does not give ϕ as a parameter.) Hence all formulas in §14.4 are specializations of formulas in §14.1.

14.5 Big q-Jacobi

Different notation See p.442, Remarks:

$$P_n(x;a,b,c,d;q) := P_n(qac^{-1}x;a,b,-ac^{-1}d;q) = {}_3\phi_2\left(\begin{matrix}q^{-n},q^{n+1}ab,qac^{-1}x\\ qa,-qac^{-1}d\end{matrix};q,q\right). \tag{105}$$

Furthermore,

$$P_n(x;a,b,c,d;q) = P_n(\lambda x;a,b,\lambda c,\lambda d;q), \tag{106}$$

$$P_n(x;a,b,c;q) = P_n(-q^{-1}c^{-1}x;a,b,-ac^{-1},1;q) \tag{107}$$

Orthogonality relation (equivalent to (14.5.2), see also [K15, (2.42), (2.41), (2.36), (2.35)]). Let $c, d > 0$ and either $a \in (-c/(qd), 1/q)$, $b \in (-d/(cq), 1/q)$ or $a/c = -\overline{b}/d \notin \mathbb{R}$. Then

$$\int_{-d}^{c} P_m(x;a,b,c,d;q) P_n(x;a,b,c,d;q) \, \frac{(qx/c, -qx/d; q)_\infty}{(qax/c, -qbx/d; q)_\infty} \, d_q x = h_n \, \delta_{m,n} \,, \tag{108}$$

where

$$\frac{h_n}{h_0} = q^{\frac{1}{2}n(n-1)} \left(\frac{q^2 a^2 d}{c} \right)^n \, \frac{1-qab}{1-q^{2n+1}ab} \, \frac{(q, qb, -qbc/d; q)_n}{(qa, qab, -qad/c; q)_n} \tag{109}$$

and

$$h_0 = (1-q)c \, \frac{(q, -d/c, -qc/d, q^2 ab; q)_\infty}{(qa, qb, -qbc/d, -qad/c; q)_\infty} \,. \tag{110}$$

Other hypergeometric representation and asymptotics

$$P_n(x;a,b,c,d;q) = \frac{(-qbd^{-1}x; q)_n}{(-q^{-n}a^{-1}cd^{-1}; q)_n} \; {}_3\phi_2 \left(\begin{matrix} q^{-n}, q^{-n}b^{-1}, cx^{-1} \\ qa, -q^{-n}b^{-1}dx^{-1} \end{matrix} ; q, q \right) \tag{111}$$

$$= (qac^{-1}x)^n \, \frac{(qb, cx^{-1}; q)_n}{(qa, -qac^{-1}d; q)_n} \; {}_3\phi_2 \left(\begin{matrix} q^{-n}, q^{-n}a^{-1}, -qbd^{-1}x \\ qb, q^{1-n}c^{-1}x \end{matrix} ; q, -q^{n+1}ac^{-1}d \right) \tag{112}$$

$$= (qac^{-1}x)^n \, \frac{(qb, q; q)_n}{(-qac^{-1}d; q)_n} \sum_{k=0}^{n} \frac{(cx^{-1}; q)_{n-k}}{(q, qa; q)_{n-k}} \, \frac{(-qbd^{-1}x; q)_k}{(qb, q; q)_k} \, (-1)^k q^{\frac{1}{2}k(k-1)} (-dx^{-1})^k. \tag{113}$$

Formula (111) follows from (105) by [GR, (III.11)] and next (112) follows by series inversion [GR, Exercise 1.4(ii)]. Formulas (111) and (113) are also given in [Ism, (18.4.28), (18.4.29)]. It follows from (112) or (113) that (see [298, (1.17)] or [Ism, (18.4.31)])

$$\lim_{n\to\infty} (qac^{-1}x)^{-n} P_n(x;a,b,c,d;q) = \frac{(cx^{-1}, -dx^{-1}; q)_\infty}{(-qac^{-1}d, qa; q)_\infty} \,, \tag{114}$$

uniformly for x in compact subsets of $\mathbb{C}\backslash\{0\}$. (Exclusion of the spectral points $x = cq^m, dq^m$ $(m = 0, 1, 2, \ldots)$, as was done in [298] and [Ism], is not necessary. However, while (114) yields 0 at these points, a more refined asymptotics at these points is given in [298] and [Ism].) For the proof of (114) use that

$$\lim_{n\to\infty} (qac^{-1}x)^{-n} P_n(x;a,b,c,d;q) = \frac{(qb, cx^{-1}; q)_n}{(qa, -qac^{-1}d; q)_n} \; {}_1\phi_1 \left(\begin{matrix} -qbd^{-1}x \\ qb \end{matrix} ; q, -dx^{-1} \right), \tag{115}$$

which can be evaluated by [GR, (II.5)]. Formula (115) follows formally from (112), and it follows rigorously, by dominated convergence, from (113).

Symmetry (see [K15, §2.5]).

$$\frac{P_n(-x;a,b,c,d;q)}{P_n(-d/(qb);a,b,c,d;q)} = P_n(x;b,a,d,c;q). \tag{116}$$

Special values

$$P_n(c/(qa);a,b,c,d;q) = 1, \tag{117}$$

$$P_n(-d/(qb);a,b,c,d;q) = \left(-\frac{ad}{bc}\right)^n \frac{(qb,-qbc/d;q)_n}{(qa,-qad/c;q)_n}, \tag{118}$$

$$P_n(c;a,b,c,d;q) = q^{\frac{1}{2}n(n+1)} \left(\frac{ad}{c}\right)^n \frac{(-qbc/d;q)_n}{(-qad/c;q)_n}, \tag{119}$$

$$P_n(-d;a,b,c,d;q) = q^{\frac{1}{2}n(n+1)}(-a)^n \frac{(qb;q)_n}{(qa;q)_n}. \tag{120}$$

Quadratic transformations (see [K15, (2.48), (2.49)] and (145)).
These express big q-Jacobi polynomials $P_m(x;a,a,1,1;q)$ in terms of little q-Jacobi polynomials (see §14.12).

$$P_{2n}(x;a,a,1,1;q) = \frac{p_n(x^2;q^{-1},a^2;q^2)}{p_n((qa)^{-2};q^{-1},a^2;q^2)}, \tag{121}$$

$$P_{2n+1}(x;a,a,1,1;q) = \frac{qax\, p_n(x^2;q,a^2;q^2)}{p_n((qa)^{-2};q,a^2;q^2)}. \tag{122}$$

Hence, by (14.12.1), [GR, Exercise 1.4(ii)] and (145),

$$P_n(x;a,a,1,1;q) = \frac{(qa^2;q^2)_n}{(qa^2;q)_n}(qax)^n \, {}_2\phi_1\left(\begin{matrix}q^{-n},q^{-n+1}\\ q^{-2n+1}a^{-2}\end{matrix};q^2,(ax)^{-2}\right) \tag{123}$$

$$= \frac{(q;q)_n}{(qa^2;q)_n}(qa)^n \sum_{k=0}^{[\frac{1}{2}n]} (-1)^k q^{k(k-1)} \frac{(qa^2;q^2)_{n-k}}{(q^2;q^2)_k\,(q;q)_{n-2k}} x^{n-2k}. \tag{124}$$

q-Chebyshev polynomials In (105), with $c=d=1$, the cases $a=b=q^{-\frac{1}{2}}$ and $a=b=q^{\frac{1}{2}}$ can be considered as q-analogues of the Chebyshev polynomials of the first and second kind, respectively (§9.8.2) because of the limit (14.5.17). The quadratic relations (121), (122) can also be specialized to these cases. The definition of the q-Chebyshev polynomials may vary by normalization and by dilation of argument. They were considered in [K4]. By [24, p.279] and (121), (122), the *Al-Salam-Ismail polynomials* $U_n(x;a,b)$ (q-dependence suppressed) in the case $a=q$ can be expressed as q-Chebyshev polynomials of the second kind:

$$U_n(x,q,b) = (q^{-3}b)^{\frac{1}{2}n} \frac{1-q^{n+1}}{1-q} P_n(b^{-\frac{1}{2}}x;q^{\frac{1}{2}},q^{\frac{1}{2}},1,1;q).$$

Similarly, by [K6, (5.4), (5.1), (5.3)] and (121), (122), Cigler's q-Chebyshev polynomials $T_n(x,s,q)$ and $U_n(x,s,q)$ can be expressed in terms of the q-Chebyshev cases of (105):

$$T_n(x,s,q) = (-s)^{\frac{1}{2}n} P_n((-qs)^{-\frac{1}{2}}x;q^{-\frac{1}{2}},q^{-\frac{1}{2}},1,1;q),$$

$$U_n(x,s,q) = (-q^{-2}s)^{\frac{1}{2}n} \frac{1-q^{n+1}}{1-q} P_n((-qs)^{-\frac{1}{2}}x;q^{\frac{1}{2}},q^{\frac{1}{2}},1,1;q).$$

Limit to Discrete q-Hermite I

$$\lim_{a\to 0} a^{-n}\, P_n(x;a,a,1,1;q) = q^n\, h_n(x;q). \tag{125}$$

Here $h_n(x;q)$ is given by (14.28.1). For the proof of (125) use (111).

Pseudo big q-Jacobi polynomials Let $a,b,c,d \in \mathbb{C}$, $z_+ > 0$, $z_- < 0$ such that $\frac{(ax,bx;q)_\infty}{(cx,dx;q)_\infty} > 0$ for $x \in z_- q^{\mathbb{Z}} \cup z_+ q^{\mathbb{Z}}$. Then $(ab)/(qcd) > 0$. Assume that $(ab)/(qcd) < 1$. Let N be the largest nonnegative integer such that $q^{2N} > (ab)/(qcd)$. Then

$$\int_{z_- q^{\mathbb{Z}} \cup z_+ q^{\mathbb{Z}}} P_m(cx;c/b,d/a,c/a;q)\, P_n(cx;c/b,d/a,c/a;q)\, \frac{(ax,bx;q)_\infty}{(cx,dx;q)_\infty}\, d_q x = h_n \delta_{m,n} \qquad (m,n = 0,1,\ldots,N), \tag{126}$$

where

$$\frac{h_n}{h_0} = (-1)^n \left(\frac{c^2}{ab}\right)^n q^{\frac{1}{2}n(n-1)} q^{2n}\, \frac{(q,qd/a,qd/b;q)_n}{(qcd/(ab),qc/a,qc/b;q)_n}\, \frac{1-qcd/(ab)}{1-q^{2n+1}cd/(ab)} \tag{127}$$

and

$$h_0 = \int_{z_- q^{\mathbb{Z}} \cup z_+ q^{\mathbb{Z}}} \frac{(ax,bx;q)_\infty}{(cx,dx;q)_\infty}\, d_q x = (1-q) z_+\, \frac{(q,a/c,a/d,b/c,b/d;q)_\infty}{(ab/(qcd);q)_\infty}\, \frac{\theta(z_-/z_+, cdz_-z_+;q)}{\theta(cz_-,dz_-,cz_+,dz_+;q)}\,. \tag{128}$$

See Groenevelt & Koelink [K12, Prop. 2.2]. Formula (128) was first given by Slater [K25, (5)] as an evaluation of a sum of two ${}_2\psi_2$ series. The same formula is given in Slater [471, (7.2.6)] and in [GR, Exercise 5.10]], but in both cases with the same slight error, see [K12, 2nd paragraph after Lemma 2.1] for correction. The theta function is given by (9). Note that

$$P_n(cx;c/b,d/a,c/a;q) = P_n(-q^{-1}ax;c/b,d/a,-a/b,1;q). \tag{129}$$

Limit relations

Pseudo big q-Jacobi $\longrightarrow$ Discrete Hermite II

$$\lim_{a\to\infty} i^n q^{\frac{1}{2}n(n-1)} P_n(q^{-1}a^{-1}ix;a,a,1,1;q) = \widetilde{h}_n(x;q). \tag{130}$$

For the proof use (124) and (180). Note that $P_n(q^{-1}a^{-1}ix;a,a,1,1;q)$ is obtained from the right-hand side of (130) by replacing a,b,c,d by $-ia^{-1}, ia^{-1}, i, -i$.

Pseudo big q-Jacobi $\longrightarrow$ Pseudo Jacobi

$$\lim_{q\uparrow 1} P_n(iq^{\frac{1}{2}(-N-1+i\nu)}x; -q^{-N-1}, -q^{-N-1}, q^{-N+i\nu-1};q) = \frac{P_n(x;\nu,N)}{P_n(-i;\nu,N)}\,. \tag{131}$$

Here the big q-Jacobi polynomial on the left-hand side equals $P_n(cx;c/b,d/a,c/a;q)$ with $a = iq^{\frac{1}{2}(N+1-i\nu)}$, $b = -iq^{\frac{1}{2}(N+1+i\nu)}$, $c = iq^{\frac{1}{2}(-N-1+i\nu)}$, $d = -iq^{\frac{1}{2}(-N-1-i\nu)}$.

14.7 Dual q-Hahn

Orthogonality relation More generally we have (14.7.2) with positive weights in any of the following cases: (i) $0 < \gamma q < 1$, $0 < \delta q < 1$; (ii) $0 < \gamma q < 1$, $\delta < 0$; (iii) $\gamma < 0$, $\delta > q^{-N}$; (iv) $\gamma > q^{-N}$, $\delta > q^{-N}$; (v) $0 < q\gamma < 1$, $\delta = 0$. This also follows by inspection of the positivity of the coefficient of $p_{n-1}(x)$ in (14.7.4). Case (v) yields Affine q-Krawtchouk in view of (14.7.13).

Symmetry

$$R_n(x;\gamma,\delta,N \mid q) = \frac{(\delta^{-1}q^{-N};q)_n}{(\gamma q;q)_n}\left(\gamma\delta q^{N+1}\right)^n R_n(\gamma^{-1}\delta^{-1}q^{-1-N}x;\delta^{-1}q^{-N-1},\gamma^{-1}q^{-N-1},N \mid q). \tag{132}$$

This follows from (14.7.1) combined with [GR, (III.11)].

14.8 Al-Salam-Chihara

Symmetry The Al-Salam-Chihara polynomials $Q_n(x;a,b \mid q)$ are symmetric in a,b.
This follows from the orthogonality relation (14.8.2) together with the value of its coefficient of x^n given in (14.8.5b).

q^{-1}-Al-Salam-Chihara

Re: (14.8.1) For $x \in \mathbb{Z}_{\geq 0}$:

$$Q_n(\tfrac{1}{2}(aq^{-x}+a^{-1}q^x);a,b \mid q^{-1}) = (-1)^n b^n q^{-\frac{1}{2}n(n-1)}\left((ab)^{-1};q\right)_n \times {}_3\phi_1\left(\begin{matrix}q^{-n},q^{-x},a^{-2}q^x\\(ab)^{-1}\end{matrix};q,q^n ab^{-1}\right) \tag{133}$$

$$= (-ab^{-1})^x\, q^{-\frac{1}{2}x(x+1)}\,\frac{(qba^{-1};q)_x}{(a^{-1}b^{-1};q)_x}\,{}_2\phi_1\left(\begin{matrix}q^{-x},a^{-2}q^x\\qba^{-1}\end{matrix};q,q^{n+1}\right) \tag{134}$$

$$= (-ab^{-1})^x\, q^{-\frac{1}{2}x(x+1)}\,\frac{(qba^{-1};q)_x}{(a^{-1}b^{-1};q)_x}\,p_x(q^n;ba^{-1},(qab)^{-1};q). \tag{135}$$

Formula (133) follows from the first identity in (14.8.1). Next (134) follows from [GR, (III.8)]. Finally (135) gives the little q-Jacobi polynomials (14.12.1). See also [79, §3].

Orthogonality

$$\sum_{x=0}^{\infty}\frac{(1-q^{2x}a^{-2})(a^{-2},(ab)^{-1};q)_x}{(1-a^{-2})(q,bqa^{-1};q)_x}\,(ba^{-1})^x q^{x^2}(Q_mQ_n)(\tfrac{1}{2}(aq^{-x}+a^{-1}q^x);a,b;q)$$
$$= \frac{(qa^{-2};q)_\infty}{(ba^{-1}q;q)_\infty}\,(q,(ab)^{-1};q)_n\,(ab)^n q^{-n^2}\,\delta_{m,n} \quad (ab>1,\ qb<a). \tag{136}$$

This follows from (29) together with (14.12.2) and the completeness of the orthogonal system of the little q-Jacobi polynomials, See also [79, §3]. An alternative proof is given in [64]. There combine (3.82) with (3.81), (3.67), (3.40).

Normalized recurrence relation

$$xp_n(x) = p_{n+1}(x) + \tfrac{1}{2}(a+b)q^{-n}p_n(x) + \tfrac{1}{4}(q^{-n}-1)(abq^{-n+1}-1)p_{n-1}(x), \tag{137}$$

where

$$Q_n(x;a,b \mid q^{-1}) = 2^n p_n(x).$$

14.10 Continuous q-Jacobi

Symmetry

$$P_n^{(\alpha,\beta)}(-x \mid q) = (-1)^n q^{\frac{1}{2}(\alpha-\beta)n}\, P_n^{(\beta,\alpha)}(x \mid q). \tag{138}$$

This follows from (97) and (14.1.19).

14.10.1 Continuous q-ultraspherical / Rogers

Re: (14.10.17)

$$C_n(\cos\theta;\beta \mid q) = \frac{(\beta^2;q)_n}{(q;q)_n}\,\beta^{-\frac{1}{2}n}\;{}_4\phi_3\left(\begin{matrix} q^{-\frac{1}{2}n}, \beta q^{\frac{1}{2}n}, \beta^{\frac{1}{2}}e^{i\theta}, \beta^{\frac{1}{2}}e^{-i\theta} \\ -\beta, \beta^{\frac{1}{2}}q^{\frac{1}{4}}, -\beta^{\frac{1}{2}}q^{\frac{1}{4}} \end{matrix}; q^{\frac{1}{2}}, q^{\frac{1}{2}}\right), \tag{139}$$

see [GR, (7.4.13), (7.4.14)].

Re: (14.10.21) (another q-difference equation). Let $C_n[e^{i\theta};\beta \mid q] := C_n(\cos\theta;\beta \mid q)$.

$$\frac{1-\beta z^2}{1-z^2}\,C_n[q^{\frac{1}{2}}z;\beta \mid q] + \frac{1-\beta z^{-2}}{1-z^{-2}}\,C_n[q^{-\frac{1}{2}}z;\beta \mid q] = (q^{-\frac{1}{2}n} + q^{\frac{1}{2}n}\beta)\,C_n[z;\beta \mid q], \tag{140}$$

see [351, (6.10)].

Re: (14.10.23) This can also be written as

$$C_n[q^{\frac{1}{2}}z;\beta \mid q] - C_n[q^{-\frac{1}{2}}z;\beta \mid q] = q^{-\frac{1}{2}n}(\beta-1)(z-z^{-1})C_{n-1}[z;q\beta \mid q]. \tag{141}$$

Two other shift relations follow from the previous two equations:

$$(\beta+1)C_n[q^{\frac{1}{2}}z;\beta \mid q] = (q^{-\frac{1}{2}n} + q^{\frac{1}{2}n}\beta)C_n[z;\beta \mid q] + q^{-\frac{1}{2}n}(\beta-1)(z-\beta z^{-1})C_{n-1}[z;q\beta \mid q], \tag{142}$$

$$(\beta+1)C_n[q^{-\frac{1}{2}}z;\beta \mid q] = (q^{-\frac{1}{2}n} + q^{\frac{1}{2}n}\beta)C_n[z;\beta \mid q] + q^{-\frac{1}{2}n}(\beta-1)(z^{-1}-\beta z)C_{n-1}[z;q\beta \mid q]. \tag{143}$$

14.11 Big q-Laguerre

Symmetry The big q-Laguerre polynomials $P_n(x;a,b;q)$ are symmetric in a,b.

This follows from (14.11.1). As a consequence, it is sufficient to give generating function (14.11.11). Then the generating function (14.1.12) will follow by symmetry in the parameters.

14.12 Little q-Jacobi

Special values (see [K15, §2.4]).

$$p_n(0;a,b;q) = 1, \tag{144}$$

$$p_n(q^{-1}b^{-1};a,b;q) = (-qb)^{-n}\, q^{-\frac{1}{2}n(n-1)}\, \frac{(qb;q)_n}{(qa;q)_n}\,, \tag{145}$$

$$p_n(1;a,b;q) = (-a)^n\, q^{\frac{1}{2}n(n+1)}\, \frac{(qb;q)_n}{(qa;q)_n}\,. \tag{146}$$

14.14 Quantum q-Krawtchouk

q-Hypergeometric representation For $n = 0, 1, \ldots, N$ (see (14.14.1) and use (7)):

$$K_n^{\mathrm{qtm}}(y;p,N;q) = {}_2\phi_1\left(\begin{matrix} q^{-n}, y \\ q^{-N} \end{matrix}; q, pq^{n+1}\right) \tag{147}$$

$$= (pyq^{N+1};q)_n\ {}_3\phi_2\left(\begin{matrix} q^{-n}, q^{-N}/y, 0 \\ q^{-N}, q^{-N-n}/(py) \end{matrix}; q, q\right). \tag{148}$$

Special values By (147) and [GR, (II.4)]:

$$K_n^{\mathrm{qtm}}(1;p.N;q) = 1, \qquad K_n^{\mathrm{qtm}}(q^{-N};p,N;q) = (pq;q)_n. \tag{149}$$

By (148) and (149) we have the self-duality

$$\frac{K_n^{\mathrm{qtm}}(q^{x-N};p,N;q)}{K_n^{\mathrm{qtm}}(q^{-N};p,N;q)} = \frac{K_x^{\mathrm{qtm}}(q^{n-N};p,N;q)}{K_x^{\mathrm{qtm}}(q^{-N};p,N;q)} \qquad (n,x \in \{0,1,\ldots,N\}). \tag{150}$$

By (149) and (150) we have also

$$K_N^{\mathrm{qtm}}(q^{-x};p,N;q) = (pq^N;q^{-1})_x \qquad (x \in \{0,1,\ldots,N\}). \tag{151}$$

Limit for $q \to 1$ to Krawtchouk (see (14.14.14) and Section 9.11):

$$\lim_{q\to 1} K_n^{\mathrm{qtm}}(1+(1-q)x;p,N;q) = K_n(x;p^{-1},N), \tag{152}$$

$$\lim_{q\to 1} K_n^{\mathrm{qtm}}(q^{-x};p,N;q) = K_n(x;p^{-1},N). \tag{153}$$

Quantum q^{-1}-Krawtchouk By (147), (149), (6) and (156) (see also p.496, second formula):

$$\frac{K_n^{\mathrm{qtm}}(y;p,N;q^{-1})}{K_n^{\mathrm{qtm}}(q^N;p,N;q^{-1})} = \frac{1}{(pq^{-1};q^{-1})_n}\ {}_2\phi_1\left(\begin{matrix} q^{-n}, y^{-1} \\ q^{-N} \end{matrix}; q, pyq^{-N}\right) \tag{154}$$

$$= K_n^{\mathrm{Aff}}(q^{-N}y;p^{-1},N;q). \tag{155}$$

Rewrite (155) as

$$K_m^{\mathrm{qtm}}(1+(1-q^{-1})qx;p^{-1},N;q^{-1}) = ((pq)^{-1};q^{-1})_n\, K_n^{\mathrm{Aff}}\Big(1+(1-q)q^{-N}\big(\tfrac{1-q^N}{1-q} - x\big);p,N;q\Big).$$

In view of (152) and (161) this tends to (70) as $q \to 1$.

The orthogonality relation (14.14.2) holds with positive weights for $q > 1$ if $p > q^{-1}$.

History The origin of the name of the quantum q-Krawtchouk polynomials is by their interpretation as matrix elements of irreducible corepresentations of (the quantized function algebra of) the quantum group $SU_q(2)$ considered with respect to its quantum subgroup $U(1)$. The orthogonality relation and dual orthogonality relation of these polynomials are an expression of the unitarity of these corepresentations. See for instance [343, Section 6].

14.16 Affine q-Krawtchouk

q-Hypergeometric representation For $n = 0, 1, \ldots, N$ (see (14.16.1)):

$$K_n^{\mathrm{Aff}}(y; p, N; q) = \frac{1}{(p^{-1}q^{-1}; q^{-1})_n}\, {}_2\phi_1\left(\begin{matrix} q^{-n}, q^{-N}y^{-1} \\ q^{-N} \end{matrix}; q, p^{-1}y\right) \tag{156}$$

$$= {}_3\phi_2\left(\begin{matrix} q^{-n}, y, 0 \\ q^{-N}, pq \end{matrix}; q, q\right). \tag{157}$$

Self-duality By (157):

$$K_n^{\mathrm{Aff}}(q^{-x}; p, N; q) = K_x^{\mathrm{Aff}}(q^{-n}; p, N; q) \qquad (n, x \in \{0, 1, \ldots, N\}). \tag{158}$$

Special values By (156) and [GR, (II.4)]:

$$K_n^{\mathrm{Aff}}(1; p, N; q) = 1, \qquad K_n^{\mathrm{Aff}}(q^{-N}; p, N; q) = \frac{1}{((pq)^{-1}; q^{-1})_n}. \tag{159}$$

By (159) and (158) we have also

$$K_N^{\mathrm{Aff}}(q^{-x}; p, N; q) = \frac{1}{((pq)^{-1}; q^{-1})_x}. \tag{160}$$

Limit for $q \to 1$ to Krawtchouk (see (14.16.14) and Section 9.11):

$$\lim_{q\to 1} K_n^{\mathrm{Aff}}(1 + (1-q)x; p, N; q) = K_n(x; 1-p, N), \tag{161}$$

$$\lim_{q\to 1} K_n^{\mathrm{Aff}}(q^{-x}; p, N; q) = K_n(x; 1-p, N). \tag{162}$$

A relation between quantum and affine q-Krawtchouk
By (147), (156), (159) and (158) we have for $x \in \{0, 1, \ldots, N\}$:

$$K_{N-n}^{\mathrm{qtm}}(q^{-x}; p^{-1}q^{-N-1}, N; q) = \frac{K_x^{\mathrm{Aff}}(q^{-n}; p, N; q)}{K_x^{\mathrm{Aff}}(q^{-N}; p, N; q)} \tag{163}$$

$$= \frac{K_n^{\mathrm{Aff}}(q^{-x}; p, N; q)}{K_N^{\mathrm{Aff}}(q^{-x}; p, N; q)}. \tag{164}$$

Formula (163) is given in [K3, formula after (12)] and [K10, (59)]. In view of (153) and (162) formula (164) has (71) as a limit case for $q \to 1$.

Affine q^{-1}-Krawtchouk By (156), (159), (6) and (147) (see also p.505, first formula):

$$\frac{K_n^{\mathrm{Aff}}(y;p,N;q^{-1})}{K_n^{\mathrm{Aff}}(q^N;p,N;q^{-1})} = {}_2\phi_1\left(\begin{matrix}q^{-n},q^{-N}y\\ q^{-N}\end{matrix};q,p^{-1}q^{n+1}\right) \tag{165}$$

$$= K_n^{\mathrm{qtm}}(q^{-N}y;p^{-1},N;q). \tag{166}$$

Formula (166) is equivalent to (155). Just as for (155), it tends after suitable substitutions to (70) as $q \to 1$.

The orthogonality relation (14.16.2) holds with positive weights for $q > 1$ if $0 < p < q^{-N}$.

History The affine q-Krwtchouk polynomials were considered by Delsarte [161, Theorem 11], [K7, (16)] in connection with certain association schemes. He called these polynomials generalized Krawtchouk polynomials. (Note that the ${}_2\phi_2$ in [K7, (16)] is in fact a ${}_3\phi_2$ with one upper parameter equal to 0.) Next Dunkl [186, Definition 2.6, Section 5.1] reformulated this as an interpretation as spherical functions on certain Chevalley groups. He called these polynomials q-Kratchouk polynomials. The current name *affine q=Kratchouk polynomials* was introduced by Stanton [488, 4.13].

14.17 Dual q-Krawtchouk

Symmetry

$$K_n(x;c,N \mid q) = c^n\, K_n(c^{-1}x;c^{-1},N \mid q). \tag{167}$$

This follows from (14.17.1) combined with [GR, (III.11)].

In particular,

$$K_n(x;-1,N \mid q) = (-1)^n\, K_n(-x;-1,N \mid q). \tag{168}$$

14.20 Little q-Laguerre / Wall

Re: (14.20.11) The right-hand side of this generating function converges for $|xt| < 1$. We can rewrite the left-hand side by use of the transformation

$${}_2\phi_1\left(\begin{matrix}0,0\\ c\end{matrix};q,z\right) = \frac{1}{(z;q)_\infty}\ {}_0\phi_1\left(\begin{matrix}-\\ c\end{matrix};q,cz\right).$$

Then we obtain:

$$(t;q)_\infty\ {}_2\phi_1\left(\begin{matrix}0,0\\ aq\end{matrix};q,xt\right) = \sum_{n=0}^{\infty} \frac{(-1)^n\, q^{\frac{1}{2}n(n-1)}}{(q;q)_n}\, p_n(x;a;q)\, t^n \qquad (|xt|<1). \tag{169}$$

Expansion of x^n

Divide both sides of (169) by $(t;q)_\infty$. Then coefficients of the same power of t on both sides must be equal. We obtain:

$$x^n = (a;q)_n \sum_{k=0}^{n} \frac{(q^{-n};q)_k}{(q;q)_k}\, q^{nk}\, p_k(x;a;q). \tag{170}$$

Quadratic transformations

Little q-Laguerre polynomials $p_n(x;a;q)$ with $a = q^{\pm\frac{1}{2}}$ are related to discrete q-Hermite I polynomials $h_n(x;q)$:

$$p_n(x^2;q^{-1};q^2) = \frac{(-1)^n q^{-n(n-1)}}{(q;q^2)_n}\, h_{2n}(x;q), \tag{171}$$

$$xp_n(x^2;q;q^2) = \frac{(-1)^n q^{-n(n-1)}}{(q^3;q^2)_n}\, h_{2n+1}(x;q). \tag{172}$$

14.21 q-Laguerre

Notation Here the q-Laguerre polynomial is denoted by $L_n^\alpha(x;q)$ instead of $L_n^{(\alpha)}(x;q)$.

Orthogonality relation

(14.21.2) can be rewritten with simplified right-hand side:

$$\int_0^\infty L_m^\alpha(x;q)\, L_n^\alpha(x;q)\, \frac{x^\alpha}{(-x;q)_\infty}\, dx = h_n\, \delta_{m,n} \qquad (\alpha > -1) \tag{173}$$

with

$$\frac{h_n}{h_0} = \frac{(q^{\alpha+1};q)_n}{(q;q)_n q^n}, \qquad h_0 = -\,\frac{(q^{-\alpha};q)_\infty}{(q;q)_\infty}\,\frac{\pi}{\sin(\pi\alpha)}\,. \tag{174}$$

The expression for h_0 (which is Askey's q-gamma evaluation [K2, (4.2)]) should be interpreted by continuity in α for $\alpha \in \mathbb{Z}_{\geq 0}$. Explicitly we can write

$$h_n = q^{-\frac{1}{2}\alpha(\alpha+1)}\, (q;q)_\alpha\, \log(q^{-1}) \qquad (\alpha \in \mathbb{Z}_{\geq 0}). \tag{175}$$

Expansion of x^n

$$x^n = q^{-\frac{1}{2}n(n+2\alpha+1)}\, (q^{\alpha+1};q)_n \sum_{k=0}^{n} \frac{(q^{-n};q)_k}{(q^{\alpha+1};q)_k}\, q^k\, L_k^\alpha(x;q). \tag{176}$$

This follows from (170) by the equality given in the Remark at the end of §14.20. Alternatively, it can be derived in the same way as (170) from the generating function (14.21.14).

Quadratic transformations

q-Laguerre polynomials $L_n^\alpha(x;q)$ with $\alpha = \pm\frac{1}{2}$ are related to discrete q-Hermite II polynomials $\widetilde{h}_n(x;q)$:

$$L_n^{-1/2}(x^2;q^2) = \frac{(-1)^n q^{2n^2-n}}{(q^2;q^2)_n}\, \widetilde{h}_{2n}(x;q), \tag{177}$$

$$xL_n^{1/2}(x^2;q^2) = \frac{(-1)^n q^{2n^2+n}}{(q^2;q^2)_n}\, \widetilde{h}_{2n+1}(x;q). \tag{178}$$

These follows from (171) and (172), respectively, by applying the equalities given in the Remarks at the end of §14.20 and §14.28.

14.27 Stieltjes-Wigert

An alternative weight function

The formula on top of p.547 should be corrected as

$$w(x) = \frac{\gamma}{\sqrt{\pi}}\, x^{-\frac{1}{2}} \exp(-\gamma^2 \ln^2 x), \quad x > 0, \quad \text{with} \quad \gamma^2 = -\frac{1}{2\ln q}\,. \tag{179}$$

For w the weight function given in [Sz, §2.7] the right-hand side of (179) equals const. $w(q^{-\frac{1}{2}}x)$. See also [DLMF, §18.27(vi)]].

14.28 Discrete q-Hermite I

History Discrete q Hermite I polynomials (not yet with this name) first occurred in Hahn [261], see there p.29, case V and the q-weight $\pi(x)$ given by the second expression on line 4 of p.30. However note that on the line on p.29 dealing with case V, one should read $k^2 = q^{-n}$ instead of $k^2 = -q^n$. Then, with the indicated substitutions, [261, (4.11), (4.12)] yield constant multiples of $h_{2n}(q^{-1}x;q)$ and $h_{2n+1}(q^{-1}x;q)$, respectively, due to the quadratic transformations (171), (172) together with (4.20.1).

14.29 Discrete q-Hermite II

Basic hypergeometric representation (see (14.29.1))

$$\widetilde{h}_n(x;q) = x^n\, {}_2\phi_1\left(\begin{matrix} q^{-n}, q^{-n+1} \\ 0 \end{matrix}; q^2, -q^2x^{-2}\right). \tag{180}$$

Standard references

[AAR] G. E. Andrews, R. Askey and R. Roy, *Special functions*, Cambridge University Press, 1999.

[DLMF] *NIST Handbook of Mathematical Functions*, Cambridge University Press, 2010; *DLMF, Digital Library of Mathematical Functions*, `http://dlmf.nist.gov`.

[GR] G. Gasper and M. Rahman, *Basic hypergeometric series*, 2nd edn., Cambridge University Press, 2004.

[HTF1] A. Erdélyi, *Higher transcendental functions, Vol. 1*, McGraw-Hill, 1953.

[Ism] M. E. H. Ismail, *Classical and quantum orthogonal polynomials in one variable*, Cambridge University Press, 2005; reprinted and corrected, 2009.

[Sz] G. Szegő, *Orthogonal polynomials*, Colloquium Publications 23, American Mathematical Society, Fourth Edition, 1975.

References from Koekoek, Lesky & Swarttouw

[24] W. A. Al-Salam and M. E. H. Ismail, *Orthogonal polynomials associated with the Rogers-Ramanujan continued fraction*, Pacific J. Math. 104 (1983), 269–283.

[46] R. Askey, *Orthogonal polynomials and special functions*, CBMS Regional Conference Series, Vol. 21, SIAM, 1975.

[51] R. Askey, *Beta integrals and the associated orthogonal polynomials*, in: *Number theory, Madras 1987*, Lecture Notes in Mathematics 1395, Springer-Verlag, 1989, pp. 84–121.

[64] R. Askey and M. E. H. Ismail, *Recurrence relations, continued fractions, and orthogonal polynomials*, Mem. Amer. Math. Soc. 49 (1984), no. 300.

[72] R. Askey and J. A. Wilson, *Some basic hypergeometric orthogonal polynomials that generalize Jacobi polynomials*, Mem. Amer. Math. Soc. 54 (1985), no. 319.

[79] N. M. Atakishiyev and A. U. Klimyk, *On q-orthogonal polynomials, dual to little and big q-Jacobi polynomials*, J. Math. Anal. Appl. 294 (2004), 246–257.

[91] W. N. Bailey, *The generating function of Jacobi polynomials*, J. London Math. Soc. 13 (1938), 8–12.

[109] F. Brafman, *Generating functions of Jacobi and related polynomials*, Proc. Amer. Math. Soc. 2 (1951), 942–949.

[146] T. S. Chihara, *An introduction to orthogonal polynomials*, Gordon and Breach, 1978; reprinted Dover Publications, 2011.

[161] Ph. Delsarte, *Association schemes and t-designs in regular semilattices*, J. Combin. Theory Ser. A 20 (1976), 230–243.

[186] C. F. Dunkl, *An addition theorem for some q-Hahn polynomials*, Monatsh. Math. 85 (1977), 5–37.

[234] G. Gasper and M. Rahman, *Positivity of the Poisson kernel for the continuous q-ultraspherical polynomials*, SIAM J. Math. Anal. 14 (1983), 409–420.

[261] W. Hahn, *Über Orthogonalpolynome, die q-Differenzengleichungen genügen*, Math. Nachr. 2 (1949), 4–34.

[281] M. E. H. Ismail, J. Letessier, G. Valent and J. Wimp, *Two families of associated Wilson polynomials*, Canad. J. Math. 42 (1990), 659–695.

[298] M. E. H. Ismail and J. A. Wilson, *Asymptotic and generating relations for the q-Jacobi and ${}_4\phi_3$ polynomials*, J. Approx. Theory 36 (1982), 43–54.

[322] T. Koornwinder, *Jacobi polynomials III. Analytic proof of the addition formula*, SIAM J. Math. Anal. 6 (1975) 533–543.

[343] T. H. Koornwinder, *Representations of the twisted $SU(2)$ quantum group and some q-hypergeometric orthogonal polynomials*, Indag. Math. 51 (1989), 97–117.

[351] T. H. Koornwinder, *The structure relation for Askey-Wilson polynomials*, J. Comput. Appl. Math. 207 (2007), 214–226; `arXiv:math/0601303v3`.

[382] P. A. Lesky, *Endliche und unendliche Systeme von kontinuierlichen klassischen Orthogonalpolynomen*, Z. Angew. Math. Mech. 76 (1996), 181–184.

[384] P. A. Lesky, *Einordnung der Polynome von Romanovski-Bessel in das Askey-Tableau*, Z. Angew. Math. Mech. 78 (1998), 646–648.

[406] J. Meixner, *Orthogonale Polynomsysteme mit einer besonderen Gestalt der erzeugenden Funktion*, J. London Math. Soc. 9 (1934), 6–13.

[463] V. Romanovski, *Sur quelques classes nouvelles de polynômes orthogonaux*, C. R. Acad. Sci. Paris 188 (1929), 1023–1025.

[471] L. J. Slater, *Generalized hypergeometric functions*, Cambridge University Press, 1966.

[485] D. Stanton, *A short proof of a generating function for Jacobi polynomials*, Proc. Amer. Math. Soc. 80 (1980), 398–400.

[488] D. Stanton, *Orthogonal polynomials and Chevalley groups*, in: *Special functions: group theoretical aspects and applications*, Reidel, 1984, pp. 87-128.

[513] J. A. Wilson, *Asymptotics for the ${}_4F_3$ polynomials*, J. Approx. Theory 66 (1991), 58–71.

Other references

[K1] R. Askey and J. Fitch, *Integral representations for Jacobi polynomials and some applications*, J. Math. Anal. Appl. 26 (1969), 411–437.

[K2] R. Askey, *Ramanujan's extensions of the gamma and beta functions*, Amer. Math. Monthly 87 (1980), 346–359.

[K3] M. N. Atakishiyev and V. A. Groza, *The quantum algebra $U_q(su_2)$ and q-Krawtchouk families of polynomials*, J. Phys. A 37 (2004), 2625–2635.

[K4] M. Atakishiyeva and N. Atakishiyev, *On discrete q-extensions of Chebyshev polynomials*, Commun. Math. Anal. 14 (2013), 1–12.

[K5] Y. Ben Cheikh and M. Gaied, *Characterization of the Dunkl-classical symmetric orthogonal polynomials*, Appl. Math. Comput. 187 (2007), 105–114.

[K6] J. Cigler, *A simple approach to q-Chebyshev polynomials*, `arXiv:1201.4703v2 [math.CO]`, 2012.

[K7] Ph. Delsarte, *Properties and applications of the recurrence* $F(i+1,k+1,n+1) = q^{k+1}F(i,k+1,n) - q^kF(i,k,n)$, SIAM J. Appl. Math. 31 (1976), 262–270.

[K8] E. Feldheim, *Relations entre les polynomes de Jacobi, Laguerre et Hermite*, Acta Math. 75 (1942), 117–138.

[K9] W. Gautschi, *On mean convergence of extended Lagrange interpolation*, J. Comput. Appl. Math. 43 (1992), 19–35.

[K10] V. X. Genest, S. Post, L. Vinet, G.-F. Yu and A. Zhedanov, *q-Rotations and Krawtchouk polynomials*, arXiv:1408.5292v2 [math-ph], 2014.

[K11] M. J. Gottlieb, *Concerning some polynomials orthogonal on a finite or enumerable set of points*, Amer. J. Math. 60 (1938), 453–458.

[K12] W. Groenevelt and E. Koelink, *The indeterminate moment problem for the q-Meixner polynomials*, J. Approx. Theory 163 (2011), 836–863.

[K13] K. Jordaan and F. Toókos, *Orthogonality and asymptotics of Pseudo-Jacobi polynomials for non-classical parameters*, J. Approx. Theory 178 (2014), 1–12.

[K14] T. H. Koornwinder, *Askey-Wilson polynomial*, Scholarpedia 7 (2012), no. 7, 7761; `http://www.scholarpedia.org/article/Askey-Wilson_polynomial`.

[K15] T. H. Koornwinder, q-Special functions, a tutorial, `arXiv:math/9403216v2 [math.CA]`, 2013.

[K16] N. N. Leonenko and N. Šuvak, *Statistical inference for student diffusion process*, Stoch. Anal. Appl. 28 (2010), 972–1002.

[K17] J. C. Mason, *Chebyshev polynomials of the second, third and fourth kinds in approximation, indefinite integration, and integral transforms*, J. Comput. Appl. Math. 49 (1993), 169–178.

[K18] J. C. Mason and D. Handscomb, *Chebyshev polynomials*, Chapman & Hall / CRC, 2002.

[K19] J. Meixner, *Umformung gewisser Reihen, deren Glieder Produkte hypergeometrischer Funktionen sind*, Deutsche Math. 6 (1942), 341–349.

[K20] N. Nielsen, *Recherches sur les polynômes d' Hermite*, Kgl. Danske Vidensk. Selsk. Math.-Fys. Medd. I.6, København, 1918.

[K21] J. Peetre, *Correspondence principle for the quantized annulus, Romanovski polynomials, and Morse potential*, J. Funct. Anal. 117 (1993), 377–400.

[K22] H. Rosengren, *Multivariable orthogonal polynomials and coupling coefficients for discrete series representations*, SIAM J. Math. Anal. 30 (1999), 233–272.

[K23] E. J. Routh, *On some properties of certain solutions of a differential equation of the second order*, Proc. London Math. Soc. 16 (1885), 245–261.

[K24] J. A. Shohat and J. D. Tamarkin, *The problem of moments*, American Mathematical Society, 1943.

[K25] L. J. Slater, *General transformations of bilateral series*, Quart. J. Math., Oxford Ser. (2) 3 (1952), 73–80.

T. H. Koornwinder, Korteweg-de Vries Institute, University of Amsterdam,
P.O. Box 94248, 1090 GE Amsterdam, The Netherlands;

email: T.H.Koornwinder@uva.nl